What Makes a Sex Life Healthy?

What science can tell us about desire, closeness, and feeling better together

Nora and Eli have sex about once a month. Last Saturday they had the house to themselves and chose pancakes and a nap. Both called it an excellent morning. If you asked either of them whether they wanted more sex, you might get the same answer. “Sometimes. We’re pretty happy.”

Alex and Jamie have sex most weeks. On Tuesday, Alex reaches for Jamie under the covers. Jamie says, “I’m exhausted.” Alex hears, “I don’t want you.” Jamie hears the silence that follows and wonders whether saying no will spoil the next day. Sometimes Jamie says yes to avoid another tense morning. If you counted only encounters, Alex and Jamie would score higher. The count would miss what it costs Jamie to decline.

Counting how often people have sex to measure intimacy is a little like counting meals to judge a dinner. You learn that someone ate. You still have to ask whether they enjoyed it, whether they felt welcome, and whether they wanted to be there.

If you’re wondering whether you have “enough” sex, start with a question the tally can’t answer. **What do I actually miss?** The World Health Organization puts well-being, safety, and freedom from coercion at the center of sexual health. Here’s what that looks like on an ordinary Tuesday. [1]

Part 1. Start with the concern you have

Ask these three questions

What do I miss? Being held? Feeling wanted? Having time together when neither of you is trying to get to sleep? “We barely have sex anymore” can mean any of those things. Say which one you mean.

What makes closeness easier? Think of one moment when touch felt welcome and one when it didn’t. Maybe Saturday morning was easier than midnight after a twelve-hour day. Maybe a kiss felt good because you knew it could end there.

Can I say no and still feel safe? A partner may feel disappointed. You should still be able to stop, change your mind, or go to sleep. If you expect pressure, threats, or retaliation, start with the printable help sheet.

> **Try this tonight.** Finish the sentence “I miss…” without using a number. If the answer is “being held,” ask for a hug. If you’re happy with things as they are, leave the tally alone.

One evening, three explanations

Maya and Rowan have a baby who regards sleep as a rumor. On the first evening they get an hour to themselves, Rowan lights a candle. Maya sees the candle and wonders how long she has to stay awake

before she can politely go to bed. They had both been looking forward to being alone. They had imagined very different uses for the hour.

Start with the **body**. Maya is tired and still figuring out what kinds of touch feel comfortable after giving birth. Then look at the **mind**. Rowan worries that the change means their attraction has faded. Finally, look at **the situation around them**. They rarely have a quiet hour. A moment of affection feels loaded because they get so few chances to have it.

Physician George Engel's biopsychosocial model asks us to look at all three levels. They can pile up. An uncomfortable encounter can make a person brace the next time. A partner can mistake that hesitation for rejection. Then both people may start avoiding the subject. A first move could be asking a clinician about pain, arranging some real rest, or saying plainly what kind of affection would feel welcome. Which one helps depends on what is happening. The same framework works for an older adult taking a new medication, a single person dating after a loss, or someone who has never especially wanted sex. [2, 3, 4, 5]

What it looks like in real life. Rowan puts the candle out. The next morning, Maya says, "I liked that you wanted some time with me. I needed sleep. Can we have coffee outside when the baby naps and see how we feel?" That is a small invitation with an easy exit. If Maya is also having pain, a conversation about coffee can coexist with a medical appointment. The relationship gets room to breathe while the pain gets the attention it deserves.



The body, mind, and circumstances shape one another.

Start with the part you can name. A change in one area can affect the others.

Find the part you need

If this sounds like you	Start here
I'm trying to figure out what counts as healthy.	Part 1. Start with the concern you have
I'm curious about the evidence.	Part 2. Read the research without the hype
I'm trying to understand a change in desire.	Part 3. Understand desire
We want different things and need words for it.	Part 4. Talk about touch
My mood, body, or circumstances have changed.	Part 5. When your body or mood changes
I'm single, starting over, or happy on my own.	Part 6. Make room for connection
I'm concerned about pain, pressure, or exposure.	Part 7. Get the right help
I want something small to try.	Part 8. Choose one next step

Part 2. Read the research without the hype

A headline can make a weekly average sound like a prescription. Here's what the studies actually asked people to do, and what happened next.

Five research snapshots you can read over coffee

Here are five findings worth remembering. The labels indicate how much weight to give each result. A strong pattern can still be an association. A well-run experiment can answer a narrow question. The labels are reading aids, not formal grades.

1. Social connection and survival

Strong pattern across studies

What they found. People with stronger social ties were more likely to be alive at follow-up. Friends, relatives, and other connections matter to this finding, too.

The study. Holt-Lunstad and colleagues pooled 148 studies with more than 300,000 participants. The studies used several measures of connection. [6]

What it can tell us. Social connection deserves attention alongside other conditions of daily life. The studies can't tell us how much any one friend, partner, or club will affect an individual's health.

What to do with it. Pick one connection you want to strengthen. Make the next step concrete. "Would you like to walk together on Wednesday?" gives someone a much easier question to answer than "How can I fix my social life?"

2. Relationship quality and physical health

Strong pattern across studies, generally small associations

What they found. People in higher-quality marriages tended to have better health on average. Across outcomes, average correlations ranged from 0.07 to 0.21 on a scale where zero means no linear link and one means a perfect positive link. The associations were modest enough that two people in equally supportive marriages could have very different health outcomes.

The study. Robles and colleagues reviewed 126 reports covering outcomes from self-rated health to survival. [7]

What it can tell us. Health and relationship quality often travel together. Work conditions, finances, and illness can also shape both. The review looked at marital quality. Its averages give little basis for judging a particular couple or assuming the same numbers apply to other relationships.

What to do with it. Ask what support would actually help this week. "Could you pick up my prescription?" and "I'd like you to listen before we solve anything" are both clearer than "I wish you'd support me."

3. Frequency and happiness

Moderate evidence about what happened when couples were asked to have more sex

What they found. In the survey data, the link between frequency and well-being leveled off around once a week for people in relationships. In a separate experiment, asking couples to have more sex brought no gain in happiness. The group given that assignment reported less desire and enjoyment.

The study. Muise and colleagues examined links between frequency and well-being in three studies totaling 30,645 participants. In a separate experiment, Loewenstein and colleagues randomly assigned some couples to try having sex more often. [8, 9]

What it can tell us. The experiment tested an assigned target in couples who agreed to take part. It leaves open what happens when people choose a change for their own reasons.

What to do with it. When you think "more," finish the sentence. More time? More comfort? More invitations you can decline? Then talk about that specific thing.

4. How people feel the next day

Moderate evidence for timing within this sample

What they found. People tended to report better well-being after days when they reported sex. Encounters rated as more pleasurable or intimate were linked with better mood the following day.

The study. Kashdan and colleagues asked 152 adults to keep diaries for three weeks. [10]

What it can tell us. A restful weekend could explain both the sex and the better mood. The diary shows which came first in this group of adults. It leaves room for other explanations and says little about how sexual difficulties and mood interact over months or years. [11]

What to do with it. Notice how *you* feel afterward. Comfortable, connected, indifferent, sore, or worried are all possible answers. Your answer may point to a conversation, a boundary, or care.

5. What intimacy means in later life

Early evidence from one study

What they found. People who put more importance on sex initially reported more depressive symptoms. Their symptoms fell faster over the period studied, and by the end their reported symptoms were lower than those of the comparison group. This was a pattern across groups of people, and individuals' experiences varied.

The study. Shen and Kwon used ten years of survey data from 1,903 partnered U.S. adults aged 57 to 85 at the start. They tracked how important sex was to participants and their depressive symptoms. [12]

What it can tell us. The study describes how two groups of partnered older adults differed over time. It leaves open why their symptoms changed differently. It did not test whether changing how much someone values sex would change their mood.

What to do with it. Ask whether the role sex plays in your life matches what you want from it. If there's a gap, name it. "I miss affection since my partner got ill" and "I'd like help with discomfort" point to different next steps.

What support looks like on an ordinary day

Support takes at least two forms. Someone can sit with you while you wait for test results. They can also ask what excites you about a class you want to take. Feeney and Collins use this distinction to explain why close ties matter in hard times and good ones. When someone shares something personal, feeling heard and understood is part of what makes the exchange intimate. "Do you want me to listen or help you figure it out?" is a useful question in either kind of conversation. [13, 14]

Can sex help with pain, heart health, or immunity?

Can sex change how pain feels? In a laboratory study, Beverly Whipple and Barry Komisaruk found that some forms of vaginal self-stimulation raised pain thresholds in women. That experiment measured how much of a test stimulus it took before someone felt pain. Treating a recurring condition such as migraine or chronic pelvic pain is a different question. In a headache-clinic survey, some people who had sex during an attack reported relief and others reported worse pain. Those mixed reports leave plenty of room for individual variation. The human study leaves the mechanism open. An older rat experiment also leaves open whether the same pain response involves opioids in humans. If sexual activity itself hurts, stop what hurts and seek care if the pain is frequent or severe. [2, 15, 16, 17]

If sexual activity brings on heart symptoms, get medical advice before resuming. The same applies if you have unstable angina or severe heart symptoms. The American Heart Association says sexual activity is probably safe once heart disease is stable. That guidance concerns how safely someone can tolerate the exertion. It does not establish that sex improves heart health over time. [18, 19]

Oxytocin is real. The magic dose is a myth. Researchers study oxytocin in social bonding, but a review found that levels measured outside the brain did not reliably match levels in the brain or spinal fluid at rest. A change in a blood sample leaves two questions open. Did the brain respond the same way? And did anyone actually get sick less often or recover faster? Those are different experiments. [20, 21]

The five-second headline check

Ask what researchers actually measured. A diary tells you what tended to happen first. A survey compares people as they were. An experiment tests a particular change. And a study about getting through a brief burst of exertion safely has little to say about whether that activity prevents disease. Read the outcome before you take the advice. [8, 9, 10, 19]

Part 3. Understand desire

Before deciding that your desire has disappeared, ask when it shows up. An invitation can land very differently on a Saturday morning than it did beside the laundry basket on Thursday night.

Four myths worth retiring

The myth	A better question
Everyone else is doing it more.	Would I want anything to change if I had never seen anyone else's numbers?
Once a week is the magic number.	What would make the intimacy I have more welcome or satisfying?
If the chemistry is real, desire always comes first.	When do I notice interest, and under what conditions?
Good lovers know what to do without asking.	What could I say that would make it easier for the other person to tell me what feels good?

"Everyone else" is usually an imaginary group composed of the loudest people we know and a few carefully selected strangers. We rarely hear about the week someone had a migraine, the Saturday a couple chose to sleep, or the relationship where both people are happy with very little sex. A number may tell you how many times something happened. It leaves you to find out whether it was wanted, comfortable, satisfying, or even the kind of connection you were looking for.

What changed when desire changed?

Imagine a song you usually enjoy playing while someone nearby is trying to sleep and the smoke alarm is chirping. You might still like the song. In that moment, other signals win your attention.

In their original model and later review, sex researchers John Bancroft and Erick Janssen describe arousal as a balance between processes that help it build and processes that inhibit it. People sometimes call these the accelerator and the brakes. Privacy, a welcome invitation, or a kind of touch you enjoy might help arousal build. Pain, fear of an interruption, or worry about what happens if you stop might get in the way. People vary in how sensitive they are to different cues. This model helps explain how the same person can respond differently on different days. It also leaves plenty of room for a straightforward "I don't want to." [3, 22]

Try a five-minute check-in with yourself. Think of one recent moment when closeness felt welcome and one when it didn't. What was different about your energy, mood, comfort, privacy, or the way you were approached? Write one sentence for each. If the answer is "I enjoyed it when I could relax because I knew a hug was just a hug," you have something specific to communicate. If you notice pain or a change that persists across situations, tell a clinician. [2, 3, 23, 24, 25]

Sometimes interest arrives late

Rosemary Basson offered a model of sexual response that takes *responsive desire* seriously. The model was developed to describe women's experiences. Its practical question is wider. Does interest sometimes turn up in response to an experience you already welcome? Some people notice a wish for sex before anything happens. Some notice it once they're already enjoying a kiss or a moment of closeness. A person may recognize both patterns in their own life. [26]

Here is the difference that matters in practice. "I'm not feeling sexual, but a kiss sounds nice" is a choice about kissing. "I'd rather read my book" is a choice about reading. Even if desire sometimes

develops after touch, you can stop the moment you want to. If you're the one asking, take the answer at face value. Treat the kiss as a complete experience if that is what you agreed to. Letting it end there makes the next invitation easier to trust.

A person can also have little interest that feels entirely right to them. An asexual person may experience little or no sexual attraction, and people who identify as asexual differ in whether they want romance, affection, or sexual activity. Attraction, desire, and behavior answer different questions. A long-standing preference that suits you is a different situation from a change that bothers you or a symptom you want evaluated. [27]

Feeling free, capable, and close

Psychologists Richard Ryan and Edward Deci argue that people tend to do better when they have room to make meaningful choices, feel capable of what they're doing, and feel connected to others. They call these needs autonomy, competence, and relatedness. Applied to intimacy, those ideas prompt three practical questions. Can I choose? Can I say what I want or learn what works? Can we still feel close when our answers differ? This is an application of a broad theory of motivation. It gives you a way to think through an evening or a relationship. [28]

Consider someone who knows they could say no and be treated with the same warmth the next day. They have room to offer a real yes, a clear no, or a smaller invitation of their own. Now imagine being worried that any feedback will be taken as an insult. Even a simple request to slow down starts to feel risky. There are ways to make honesty easier. Ask for one change at a time. If you hear a request, say what you heard before explaining yourself. And when someone says, "I'd prefer that we stop," let the conversation pause with the activity.

Chemistry can give you a start. Learning carries you further.

Psychologist Jessica Maxwell and colleagues compared two ideas about sex. One puts more weight on finding a naturally compatible partner. The other emphasizes what people can learn together. Across six studies that included surveys, diaries, and an experiment, stronger growth beliefs generally went with greater sexual and relationship satisfaction. Compatibility beliefs had a more mixed pattern. This research offers a question to ask after a clumsy night, "What could we learn from that?" It does not establish that changing your beliefs will reliably improve your sex life. [29]

Try it with a specific problem. "When we only bring this up in bed, I feel put on the spot. Could we talk on the walk tomorrow?" Or, "We both like being close, and our evenings are a mess. Is there a better time?" A learning mindset leaves room for two people to conclude that they want different things. It also leaves room to say that a boundary is settled. The effort goes into listening and finding choices people can genuinely agree to.

Why "not tonight" can sound like "I'm leaving"

Attachment theory offers one way to think about why a refusal can sting. When you already feel unsure about a relationship, "not tonight" may sound like "something is wrong between us." Fatigue, pain, a recent argument, or simply wanting something else could explain the answer. If the worry lingers, ask about it later, when nobody is deciding about sex. One evening gives you little basis for labeling either person's attachment style. [30]

Part 4. Talk about touch

Start this conversation while you're both dressed and nobody's waiting for an answer. A walk or ten minutes at the kitchen table gives both people more room to speak.

> **How to use these tools.** The scripts and short exercises are optional ways to apply ideas from relationship research. The studies cited here did not test these exact versions. Use what fits, skip what doesn't, and keep stopping easy. If saying no feels unsafe, go to the help sheet.

When you want different amounts of sex

One person reaches for a kiss. The other stiffens, anticipating a decision they don't want to make. The first person feels rejected and pulls away. After a while, both stop reaching for each other, even though both miss being close.

Differences in desire can show up in frequency, timing, or the kind of touch someone wants. A person may enjoy affection on Saturday morning and feel entirely unavailable after a late shift. Some people notice desire before touching. For others it sometimes develops during contact they already want and feel free to stop. That possibility depends on genuine willingness. If someone is unsure or wants to stop, let that answer stand. [32, 33]

If you and a partner want different things, make it easy to speak honestly. Pick a calm time outside a sexual encounter. Name one thing you miss or enjoy and ask one open question. For example, "I miss feeling close to you. What kind of closeness feels welcome lately?" A 93-study meta-analysis found that the perceived quality of sexual communication was more strongly associated with satisfaction than how often people talked about sex. The script is a way to start a conversation. Its value depends on whether both people can answer honestly and be heard. [34]

Try ending the conversation with one optional, specific idea both people welcome. That might be a walk together, kissing with no expectation of going further, or setting aside time to talk again. If touch often feels like an obligation, agree on an option with a clear stopping point and respect it. A response like "Thanks for telling me. We can stop here" makes that choice credible. At the next check-in, ask what felt comfortable or connecting and what each person would change. [1, 32]

Why you say yes can matter as much as whether you say it. Muise and Impett use the term sexual communal strength for a person's motivation to respond to a partner's sexual needs. In their studies, partners who felt that care reported greater relationship satisfaction and commitment. A related line of research asks what someone hopes to gain or avoid by saying yes. Saying yes because you look forward to pleasure or closeness can feel different from saying yes because you fear conflict or disappointment. Motives can overlap. You can enjoy doing something that makes your partner happy and still ask yourself, "Would I feel free to decline if I wanted to?" [31, 32, 35]

A conversation you can borrow

1. **Ask for a time to talk.** "Could we talk this weekend about how intimacy has been feeling for each of us? I'd like to understand what's been going on."
2. **Describe one moment.** "Last night I wanted to cuddle, but I worried it would turn into a decision about sex." Or, "When we went a while without touching, I started wondering if you still liked being close to me."

3. **Check what you heard.** “So you’d enjoy more affection if you knew it could end there. Did I get that right?”
4. **Agree on one small change, if you both want one.** “Would you like to cuddle while we watch a show and leave it at that?” Give it a try only if both people are comfortable. Later, ask whether the agreement was easy to trust.

If you hear “I’m not in the mood,” an answer as simple as “Okay. Would you like some company, or would you rather have space?” keeps the choice open. Accept either answer. If a mismatch keeps bothering you and both partners feel safe speaking freely, a qualified clinician or therapist with training in sexual health can help you sort out what each person wants and what gets in the way. Sometimes two people discover a workable arrangement. Sometimes they learn that their preferences differ in a way that matters to both of them. [32, 33, 34]

When a no lands hard

It can sting to be turned down, especially when you were asking for closeness as much as sex. The feeling is worth understanding. The other person gets to keep their answer. Try asking yourself what story arrived with the sting. “They’re tired” can become “They don’t find me attractive” in the time it takes someone to turn off a lamp. If the worry keeps coming back, bring it up later as a worry you want to share, with room for the other person’s experience.

In studies by James Kim and colleagues, people who saw a partner’s refusal as reassuring reported greater sexual and relationship satisfaction. Those who saw it as hostile reported lower satisfaction. If warmth feels genuine and safe, “I’m glad you asked. I’m going to sleep now” is a full answer. A simple no is enough, too. [36]

> You can care about your partner and still say no.

If you did the asking: “Okay. I’m a little disappointed, and I can handle that. Good night.” If you want to reconnect, ask another time. **If you’re declining:** “I’d like to hold hands for a minute,” or “I’d like some space.” Offer only what you actually want. **If the moment already went badly:** “I got quiet after you said no and I can see how that might have felt. I’m going to take care of my own disappointment. You get to say what you want.”

The ten-minute check-in

Conversation tool. This ten-minute format has not been directly tested in the studies cited here.

Use this when both people feel safe and want to talk. Put the phones down. Set a timer if it helps the conversation feel finite. A kitchen table or a walk can work better than a bed because neither person is being asked to decide about sex while they speak.

1. **Minutes 1–2. Say one thing you liked.** “I liked it when you put your hand on my shoulder after I got home.” Choose something real and small. “You finally did the dishes” will probably send the discussion in a different direction.
2. **Minutes 3–5. Name one friction point each.** “When we’re both exhausted, I find it hard to switch gears.” Listen for the particular moment. Repeat back what you heard before presenting your explanation.

3. **Minutes 6–8. Pick one thing to try.** “Could we take a walk after dinner on Friday?” “Would you like a hug hello, with no expectation that it becomes anything else?” An invitation can be declined. If there’s no shared idea today, agree to revisit it or let the matter rest.
4. **Minutes 9–10. Close the loop.** “What did I miss?” “When should we check in again?” Next time, ask how the small change felt to each of you. Did it make the week easier? Was it welcome? Would you repeat it?

If one partner tends to do all the scheduling, choose a fair way to share that work. If the conversation circles back to old grievances, pick a narrower subject or talk at a calmer time. If a person expects retaliation for an honest answer, skip the joint exercise and use private support.

A yes, no, maybe list

Conversation tool. Use it to name preferences. An entry on a list does not give consent later.

Write three private columns. Start with ordinary possibilities: holding hands in public, sitting close on the couch, kissing, talking about a fantasy, having more time alone. You can add whatever feels relevant, including conditions such as “only when there’s time,” “ask first,” or “no photos.” Share only the parts you want someone else to see.

A **yes** means the idea sounds welcome in some circumstances. Ask again in the moment. A **no** marks something you want left alone. A **maybe** might mean you’d enjoy talking about it or that timing, privacy, or trust would affect your answer. Agree on how to ask before assuming anything. If you compare lists, start with one place where you already overlap. It might be “we both want more time to talk.” Leave the unmatched items there unless both people want to discuss them. [32, 33]

This can also work when you’re single. Your list can include what you want to know before a date comes home with you, who you want to tell about a new relationship, or what type of touch you prefer to keep private. The point is to put your own preferences into words before you feel hurried.

When sex starts to feel like a performance

It can be hard to notice pleasure when you’re busy checking whether your body is doing what you think it should. A person who loses an erection may spend the next encounter anticipating it. Someone who usually takes longer to orgasm may start rushing because they’re worried about a partner getting tired. And someone who expects pain may find it hard to pay attention to anything else.

If you both want to stay close, consider saying what would make the moment easier. “Can we slow down?” “I’d like to stop here.” “I’m happy just lying beside you.” You can talk about pleasurable things you already know you like and make room to change your mind. If sexual activity has become uncomfortable, a comfortable alternative can be enough for that moment. Recurring pain or erection difficulty deserves care even when there are also worries about performance. [2, 25, 32]

Step out of the commentary booth

Some clinicians call it *spectatoring* when you start watching and grading yourself from the sidelines. Your attention goes to how your body looks, whether an erection will last, or how much time is passing. Meanwhile, your partner may be having a parallel monologue about something else entirely. It can be hard to enjoy a moment when you’re also reviewing the footage.

See **What has research support?** for what a clinical mindfulness program tested.

If you catch yourself grading the moment, try noticing one sensation you freely chose to experience, such as the warmth of a hand or the pressure of the couch beneath you. You can also ask for a change or stop. If focusing on your body makes you feel less safe, look around the room or talk with someone you trust. You get to decide whether any exercise is right for you.

Five minutes of touch with nowhere to go

Adapted from sensate focus. The clinical studies tested structured programs. They did not test this short exercise.

Choose this exercise only if both people want touch and feel safe stopping. The clinical research is summarized in Part 5.

- **Agree on the entire plan before you start.** You might choose two minutes of touching a hand or forearm over a sleeve, followed by two minutes with the roles switched if you both still want to. Keep it nonsexual and stay within the area you agreed on.
- **Pick plain words.** "Lighter," "slower," "stay there," and "stop" all count. Either person can end the exercise at any time. The person giving touch can pause or decline too.
- **Notice rather than perform.** Does the touch feel warm, cool, neutral, pleasant, or unwelcome? Every answer tells you something. If neither person likes it, a conversation or sitting near each other might suit you better.
- **End when you said you would.** Ask "Anything you'd keep or change?" Then go on with your day. Leaving it there shows that the agreement was real.

There are less formal options. A hand on a shoulder after asking, a few minutes sitting close while watching a show, or leaning against each other on a bench can be enough. If touch is tiring, uncomfortable, or unwelcome, share a cup of tea, a favorite song, or a walk. The useful choice is one that people actually want to make together.

Part 5. When your body or mood changes

If sex is one of several things you've stopped enjoying, look at the rest of the week. If the change began with a new medication, a new baby, or recurring pain, start there.

When your body or your life changes

If desire, arousal, or comfort has changed, describe the change precisely. A loss of interest, pain, difficulty with erections, and difficulty reaching orgasm can feel similar from the outside and have different causes. Ask when the change began, whether it happens in every situation or only in some, and what else changed around that time. Stress, mood, a new medicine, pregnancy or postpartum changes, menopause, and health conditions can all be relevant. A brief private note can make a medical conversation easier if the change persists or bothers you. [2, 23, 24, 25]

Different stages, different questions

After having a baby. Recovery, exhaustion, feeding, and finding private time can all shape how you feel about sex. If you want intimacy but aren't ready for a particular activity, say what you would welcome now. A partner can also ask what would help you feel rested or supported during the day. If discomfort persists, bring it up with your clinician. If avoiding pregnancy matters to you, ask about postpartum contraception even if your period hasn't returned. [2, 5]

Around menopause. If penetration has become uncomfortable and dryness seems to be part of it, ask about options such as a lubricant, a vaginal moisturizer, or whether local treatment is appropriate for you. If pain keeps happening, describe where it occurs and when it started so a clinician can consider other causes. A useful opening is "I still want intimacy, but I've started avoiding it because it hurts." [2, 37]

When you're living with illness, disability, or fatigue. Start with the part you can name. "Mornings are when I have energy," "That position strains my hip," or "I'm worried because I felt breathless last time." The first two may point to changing timing or choosing more comfortable kinds of touch. A new symptom is something to raise with your clinician, especially if you have a heart condition. You can ask, "Given my symptoms and treatment, what kinds of sexual activity are safe for me right now?" [2, 18]

When a medication or mood change coincides with a sexual change. You may feel relief that treatment is helping your mood and frustration that sex feels different. Tell your prescriber both parts. Include whether the change began before or after the medication, what you've noticed, and how much it bothers you. Together you can weigh what's helping and what options are appropriate. [23, 24]

When you've lost interest in more than sex

Picture someone who's stopped answering friends' messages, stopped putting on music while making dinner, and also lost interest in sex. They might first notice the sexual change because a partner mentions it. The rest of the picture matters. Compare that with someone who still enjoys friends, hobbies, and affection but has started avoiding penetration because it hurts. Both may say "My sex drive is gone," yet the first question to ask in each case is different.

A review of prospective studies found that depression predicted later sexual difficulties and that sexual difficulties also predicted later depression. That pattern emerged over longer periods than the next-day diary study, and the studies measured different things. It supports looking in both directions. It can also be a relief to have language for both problems at an appointment. "I've lost interest in a lot of things lately, and my sex drive has changed. Could we look at both?" [10, 11]

Stress, shame, and the days your mind is elsewhere

Stress can crowd the room. Imagine trying to switch from finishing a difficult work email to wanting intimacy in the time it takes to brush your teeth. Some people welcome sex as a break from stress. Some find that their attention stays with the problem. The dual control model gives you a way to ask what else is competing for attention without making a verdict about anyone's attraction. You might need sleep, more privacy, help with an unfinished task, or time alone. [3]

Anxiety can turn a moment into a forecast. "What if it hurts again?" is information worth taking seriously. "What if I disappoint them?" may tell you that the stakes feel too high. Try separating the prediction from the choice. "I'm worried I'll be uncomfortable. I'd like to keep my clothes on and just sit with you." A partner can help by taking that request literally. If anxiety is interfering in several parts of life, bring the bigger pattern to a clinician. [2, 24, 38, 39]

Shame has a way of speaking in verdicts. "My body is wrong." "I should be better at this." Try responding as you would to a friend who said the same thing. First name the moment, "I'm feeling exposed and worried about being judged." Then choose one kind sentence you can believe, "I can ask for what would make me comfortable." If kind language feels out of reach, use neutral language, "I'm allowed to slow down." Research links self-compassion with fewer body image concerns. This short practice is one way to use that idea, with room for professional support if shame or body distress persists. [40]

Sleep and alcohol are worth noticing. In a pilot diary study, longer sleep was linked to greater desire the following day in the women who took part. That finding makes sleep a reasonable thing to consider when your sex life feels different, without implying a predictable effect for everyone. Alcohol can alter sexual function and can affect someone's ability to consent. If either person is unable to give clear, voluntary consent, pause. If you often rely on a drink to make closeness tolerable, consider what you would need to feel comfortable while sober and discuss it at a time when you can both think clearly. [24, 32, 41]

Past experiences can return in the present. A sound, a position, or a feeling of being unable to move can bring up a memory or a strong reaction. You can stop, get dressed, sit somewhere else, or ask someone to leave. You can say "I need a pause" without giving your history. A partner can make space and ask whether you want company. A trauma-informed clinician can help you explore options if reactions recur or distress you. Choose the pace and the details you share. [32, 42]

What has research support?

Treatments studied for sexual concerns

Randomized trial in a specific clinical group. In Brotto's trial of women with sexual interest/arousal disorder, desire and arousal improved in both the mindfulness-based therapy and supportive sex education and therapy groups. Sexual distress improved more in the mindfulness group. Both were structured programs. [38]

Early evidence for one sensate-focus program. A trial in Chinese heterosexual couples tested an online program and reported benefits on some outcomes. That result applies to the program and people studied. [43]

Review of psychological treatments. Some interventions helped with some sexual difficulties. Results varied by treatment and concern. Ask a therapist what they would use for yours and why. [44]

Treatments for depression or anxiety

Cognitive behavioral therapy, or CBT, is an evidence-based option for depression and conditions such as generalized anxiety disorder. Ask a clinician what fits your symptoms. If you take medication, you can bring up sexual side effects alongside what the medication is helping with. [23, 45, 46]

Where the at-home tools fit

The short touch exercise in Part 4 borrows an idea from sensate focus. The self-compassion exercise above draws on research about body image. The cited studies did not test these versions as treatments for sexual concerns. You can use them to notice what feels comfortable or to put a concern into words. [40, 43]

> **When to see a professional about mood.** If low mood, anxiety, or loss of interest is interfering with sleep, work, or relationships, ask for help. Mention changes that last about two weeks, get worse, or interfere with daily life. If you are in a suicidal crisis or emotional distress, use the crisis resources in the printable help sheet. In immediate danger, call your local emergency number. [39, 45, 46, 47]

Part 6. Make room for connection

A quiet apartment can feel like freedom on Friday and loneliness on Sunday. Ask what kind of company you want before deciding where to look for it.

If you're single, starting over, or happy on your own

Sam has been single for a year. On Fridays, the apartment feels wonderfully quiet. On Sundays, the same quiet feels like there's nobody to tell about a ridiculous thing that happened at the grocery store. Sam downloads a dating app, then realizes the thing they wanted most that day was a standing lunch date with a friend. They may also want a romantic or sexual connection. These are separate wants, and both are allowed to matter.

The social-connection snapshot in Part 2 includes friendships and community ties. Start with the kind of connection you actually miss.

Get specific about the loneliness. Do you miss company, being understood, touch, sex, or a person you've lost? Company might mean a regular walk or volunteering somewhere you see the same people. Being understood might mean talking with an old friend or finding a group where some part of your experience is familiar. Missing a particular person can feel different from either. A grief group or a conversation with someone who knew them may be a place to start.

Ask for a concrete kind of contact. "I'm usually free Sunday morning. Want to get coffee twice this month?" is easier to answer than "We should hang out." If someone declines, you learned something about that invitation. Try another date or another person if you want. A partner can be one part of a network of care. A network can also be built without dating.

If you want sex or romance, you can be clear about that too. Decide what pace you want, where you want to meet, and how you'd like to talk about protection and privacy. For LGBTQ+ readers, privacy can include who knows about a relationship or what language you use about your body. Discuss this before anyone assumes they can share something. A clinician can ask about the actual activities and exposure involved in testing and prevention. Identity labels alone give them limited information about which tests you need. [48, 49]

If access is part of the problem, say so. Illness or disability can affect energy, transport, privacy, or what feels comfortable. You may need a venue with a quiet room, time when pain is lower, or a date

who understands that plans may change. At a medical visit you can ask which adaptations are safe with your condition or treatment. In everyday life you can say "Here's what would make this possible for me," and see how the other person responds.

If you have little or no interest in sex, make room for the connections you do want. An affectionate friendship, a romantic relationship with little or no sex, or a life built around chosen family can be meaningful on its own terms. Similarly, wanting sex after widowhood or later in life can fit the life you are building. A national study of older U.S. adults documented a wide range of sexual activity and concerns. It gives clinicians a reason to make space for the subject and readers room to bring it up at any age. [12, 27, 50]

A social life that fits the time you have

A familiar piece of advice is "Join something." It can be good advice and still leave you looking at a list of events you cannot afford, cannot get to, or cannot face after work. Instead, start with what kind of contact you can realistically repeat. A weekly call on the way home is one answer. A library group that meets every other Saturday is another. The point of repetition is that each contact need not carry the whole burden of becoming friends.

Think about three levels. **A person you could text today.** This might be someone who already likes hearing from you, even if it has been a while. **A place where you might see the same people again.** It could be an online group, a class, a community gathering, or a regular shift helping somewhere. **A person you could ask for practical help.** Maybe a neighbor can pick something up while you are sick, or a sibling can help you work through a form. Those are different forms of connection.

If you have low energy, an accessible option counts. Choose a gathering where you can leave early. Ask if there is a quieter space. Invite someone who is comfortable visiting at home if that feels safe. If it takes an hour to get there, the plan may work better every other week. A useful routine is one that survives contact with your actual calendar.

At the end of a few weeks, ask what gave you pleasure or relief. Did you enjoy talking with someone particular? Did you prefer doing something alongside people with less conversation? Would you like to invite one person for coffee, or are you happy with the group as it is? You get to decide whether you want to go deeper. Casual connections can have value without becoming something else.

Part 7. Get the right help

You don't have to settle what a symptom or a partner's behavior means before asking for help. The printable help sheet puts the first steps in one place.

When saying no feels unsafe

Consent is voluntary, specific to the activity, and can be withdrawn at any time. If someone goes quiet, freezes, or pulls away, stop and check in. If saying no could bring threats, intimidation, or retaliation, seek confidential support from a sexual assault or domestic violence service or a clinician when you can safely do so. After a recent assault, medical care can address injuries, pregnancy and STI concerns, and time-sensitive prevention. If there may have been HIV exposure, seek care promptly. HIV post-

exposure prophylaxis should be started as soon as possible when indicated, within 72 hours. [1, 32, 51]

One useful distinction is what happens *after* a boundary is stated. A partner can feel disappointed and still respond with care. Repeatedly arguing after a refusal, threatening consequences, or making affection conditional on sex can change whether a person feels free to choose the next time. If you recognize that pattern, consider reaching out privately from a device you can safely use. An advocate can help you think through options. When there is abuse, individual support and safety planning are a better starting point than joint counseling. [32, 52]

A safer-sex conversation you can have before things get intimate

Bring it up when you have enough privacy and time to think. “Before we have sex, I’d like us to talk about testing and protection. When were you last tested, and what was included?” Share what you know about your own testing, talk about barriers you want to use, and discuss contraception if pregnancy is possible and you want to avoid it. You can agree to wait or choose something you both feel comfortable with while you sort things out. [48, 49]

Tests vary by infection and by the kind of contact you’ve had. Many STIs cause no noticeable symptoms. A clinic can help you decide what to test for, when to test, and whether a throat or rectal swab is relevant. If a possible HIV exposure was recent, seek prompt medical advice about prevention rather than waiting for a routine test. [48, 49, 51]

What to bring to a health appointment

You can open with “I’d like to talk privately about a change in my sexual health.” Bring three details if you can.

1. **What changed and when.** Name the concern in ordinary words and say whether it is occasional, recurring, or persistent.
2. **What else was happening.** Bring relevant symptoms, major life changes, and a list of medicines and supplements. Mention whether the concern varies by situation.
3. **What you want help with.** You might want sex to feel comfortable, understand a change in desire, discuss testing or prevention, or find ways to feel close during illness or recovery.

You can bring a written note or start with a single sentence if you feel embarrassed. If the first answer leaves you unsure what to do, ask “What should I try next, and when should I come back if it’s still happening?” [48]

If an appointment is hard to arrange

Start with the kind of care you can reach. A primary care office can review a medicine or a new symptom. A sexual health clinic can help with testing and prevention. A licensed mental health professional can help you sort out distress, anxiety, or a recurring pattern. Ask what they work with and what a first visit looks like. If a specialist has a long wait, ask the referring office what you can do while you wait and which symptoms should move the appointment up. [23, 39, 48, 49]

Before you hang up or leave, write down one next step. A date, a test result to look for, or a number to call is easier to remember than an entire conversation. You can ask someone you trust to help with

the practical part, such as transport or making time for a call, without sharing more detail than you want to.

Part 8. Choose one next step

Pick one move that fits your situation. A question asked at the right time is enough for one week.

Four questions you can answer on paper

1. **When did I last feel close to someone in a way I enjoyed?** Include friendship or family as well as sexual or romantic contact. What made it possible? Who listened? What was different that day?
2. **Is there something I'm saying yes to because declining feels costly?** If the cost is a brief, ordinary disappointment, you can make room for it. If the cost is retaliation, threats, or pressure, reach for confidential support.
3. **Is the change specific to sex, or have I lost interest in other things too?** Note what changed and roughly when. If symptoms persist or interfere with life, bring those details to a health visit.
4. **What is one step I can take without asking someone else to change their mind?** Make an appointment, invite a friend for coffee, ask for a conversation, make room to rest, or decide the situation suits you.

Write an answer only if it is safe to keep it private. You can do the exercise in your head. Some answers may be uncertain. "I don't know whether I want more sex or more affection" is useful information. It tells you what question to ask next.

A week with one small aim

Early in the week, name the concern. "I miss laughing together" is more specific than "Our relationship is broken." "I feel sore after sex" is a symptom you can bring to an appointment. Give the question a size you can work with.

Midweek, make one request. Invite a friend to walk. Ask your partner if they'd like a ten-minute check-in. Make a health appointment. If the problem is urgent or safety is in doubt, use the help sheet now.

At the end of the week, decide what to keep. Did the change give you more ease, more choice, or a clearer question? If it helped, repeat it. If it didn't, change the plan. You don't owe the plan another week.

What we still don't know

This article draws on selected research reviews, diary and follow-up studies, a randomized experiment, and clinical guidance. Each contributes a different piece. Reviews show whether patterns recur across studies. Diaries can show what tends to happen first. The experiment tests what happens when couples are asked to increase frequency. Clinical guidance helps answer specific questions about safety and care. Several studies focused on married people, newlyweds, or partnered older adults, so their averages may travel poorly across ages, cultures, sexual orientations, gender

identities, and relationship arrangements. Researchers also differ in what they count as sex and what they measure afterward. The conversation starters and checklist translate the research into things a reader can try or ask about. The cited studies did not directly test those scripts. [8, 9, 10, 12, 14, 19, 32, 53]

The next morning

Nora and Eli's pancakes can stay pancakes. Both are comfortable with how often they have sex. They can ask for a change if that ever stops being true.

At Alex and Jamie's breakfast table, a different conversation is overdue. If Jamie feels safe enough to speak, they might say, "Sometimes I say yes because I'm worried about how things will feel between us tomorrow." Alex might feel stung hearing that. What matters next is whether Alex listens, takes responsibility for their own response, and gives Jamie room to choose. If Jamie expects threats or retaliation, private support is the place to begin.

Perhaps the next evening ends with a hug. Perhaps it ends with two people reading different books. If one of them reaches over again next week, the change worth noticing is whether either person can answer honestly.

Optional deep dive. When an awkward moment sticks

Repair the small thing while it is still small

In a close relationship, a clumsy comment can stay in circulation for years. Somebody jokes about how often you initiate. Somebody else laughs and remembers it every time they think about asking. The person who made the joke may have forgotten it before the plates were cleared. The person who heard it may have adjusted their behavior around it.

Try a repair that fits the size of the problem. “I said something about how long it takes you to get ready. It sounded like I was grading you. I can see how that could make you want less attention from me. I’m going to leave those jokes alone.” If the other person says it bothered them for another reason, listen. A repair can be brief and still make room for facts you did not know.

Another common repair concerns the timing of a request. “I brought it up while we were already in bed, and it put you on the spot. Could we talk somewhere else next time?” You can name your part without asking the other person to confirm that everything is fine. Give them room to have a different experience. A sincere apology may start the process. A changed pattern is what the other person gets to observe.

What does change look like? If you said you would stop making a joke, stop making it. If you agreed that a hug can end at a hug, practice ending there. If you promised to listen before making suggestions, ask what kind of response the person wants. These are small things someone can notice. They are easier to trust than “I’m going to be a completely different person.”

Sometimes repair asks for patience. Suppose you used to take every decline personally. Now you respond differently, but your partner still braces. They may be waiting to see whether the change will last. You can allow a slow update. Respecting someone’s limits counts even when their response does not immediately reassure you. If either person is afraid of what an honest conversation might bring, a private conversation with a clinician or advocate is a better first step than a planned couples exercise. [32, 52]

Printable help sheet

What's happening	Where to start
Sexual activity repeatedly hurts, or discomfort is severe.	Stop the activity that hurts. Ask a clinician about the pain and describe when and where you feel it.
Desire, erections, or orgasm changed after starting a medication.	Contact the prescriber to review timing and options before making changes yourself.
Low mood or loss of enjoyment is affecting several parts of your life.	Talk with a health or mental health professional. Mention sleep, energy, and sexual changes alongside mood.
You have unstable or severe heart symptoms, or sexual activity brings symptoms on.	Get medical advice before resuming sexual activity.
You fear threats, punishment, or pressure if you decline.	Seek confidential support when you can do so safely. A clinician or advocate can help you think through options.
There may have been a recent HIV exposure.	Seek urgent medical care. HIV PEP, when indicated, should start as soon as possible and within 72 hours.

U.S. support. If someone is in immediate danger, call **911**. For suicidal crisis or emotional distress, call 988, text 988, or use https://988lifeline.org/chat/. For domestic violence support, call **800-799-7233** or visit The National Domestic Violence Hotline. For sexual assault support, call **800-656-4673** or visit RAINN. Use a device you can safely access. Outside the U.S., use your local emergency number or contact a local crisis, domestic violence, or sexual assault service. [47, 54]

Clinical sources for this sheet. Pain and sexual symptoms [2, 23, 25]. Heart symptoms [18]. Mood [39]. Assault and HIV exposure [51]. Relationship safety [52].

Research and practical resources

1. World Health Organization. Sexual health (https://www.who.int/teams/sexual-and-reproductive-health-and-research-%28srh%29/areas-of-work/sexual-health). Accessed November 9, 2026.
2. American College of Obstetricians and Gynecologists. When sex is painful (https://www.acog.org/womens-health/faqs/when-sex-is-painful). Accessed November 9, 2026.
3. Janssen, E., & Bancroft, J. (2023). The dual control model of sexual response: A scoping review, 2009–2022. *The Journal of Sex Research, 60*(7), 948–968. https://doi.org/10.1080/00224499.2023.2219247
4. Engel, G. L. (1977). The need for a new medical model: A challenge for biomedicine. *Science, 196*(4286), 129–136. https://doi.org/10.1126/science.847460
5. NHS. Sex and contraception after birth (https://www.nhs.uk/baby/support-and-services/sex-and-contraception-after-birth/). Accessed November 9, 2026.
6. Holt-Lunstad, J., Smith, T. B., & Layton, J. B. (2010). Social relationships and mortality risk: A meta-analytic review. *PLOS Medicine, 7*(7), e1000316. https://doi.org/10.1371/journal.pmed.1000316
7. Robles, T. F., Slatcher, R. B., Trombello, J. M., & McGinn, M. M. (2014). Marital quality and health: A meta-analytic review. *Psychological Bulletin, 140*(1), 140–187. https://doi.org/10.1037/a0031859
8. Muise, A., Schimmack, U., & Impett, E. A. (2016). Sexual frequency predicts greater well-being, but more is not always better. *Social Psychological and Personality Science, 7*(4), 295–302. https://doi.org/10.1177/1948550615616462
9. Loewenstein, G., Krishnamurti, T., Kopsic, J., & McDonald, D. (2015). Does increased sexual frequency enhance happiness? *Journal of Economic Behavior & Organization, 116*, 206–218. https://doi.org/10.1016/j.jebo.2015.04.021
10. Kashdan, T. B., Goodman, F. R., Stiksma, M., Milius, C. R., & McKnight, P. E. (2018). Sexuality leads to boosts in mood and meaning in life with no evidence for the reverse direction: A daily diary investigation. *Emotion, 18*(4), 563–576. https://doi.org/10.1037/emo0000324
11. Atlantis, E., & Sullivan, T. (2012). Bidirectional association between depression and sexual dysfunction: A systematic review and meta-analysis. *The Journal of Sexual Medicine, 9*(6), 1497–1507. https://doi.org/10.1111/j.1743-6109.2012.02709.x
12. Shen, S., & Kwon, S. (2026). Motivated or misaligned? The importance of sex and mental health among partnered older adults. *Aging & Mental Health*. Advance online publication, June 4, 2026. https://doi.org/10.1080/13607863.2026.2681946
13. Feeney, B. C., & Collins, N. L. (2015). A new look at social support: A theoretical perspective on thriving through relationships. *Personality and Social Psychology Review, 19*(2), 113–147. https://doi.org/10.1177/1088868314544222
14. Laurenceau, J.-P., Barrett, L. F., & Pietromonaco, P. R. (1998). Intimacy as an interpersonal process: The importance of self-disclosure, partner disclosure, and perceived partner responsiveness in interpersonal exchanges. *Journal of Personality and Social Psychology, 74*(5), 1238–1251. https://doi.org/10.1037/0022-3514.74.5.1238

15. Whipple, B., & Komisaruk, B. R. (1985). Elevation of pain threshold by vaginal stimulation in women. *Pain, 21*(4), 357–367. https://doi.org/10.1016/0304-3959%2885%2990164-2
16. Hambach, A., Evers, S., Summ, O., Husstedt, I. W., & Frese, A. (2013). The impact of sexual activity on idiopathic headaches: An observational study. *Cephalalgia, 33*(6), 384–389. https://doi.org/10.1177/0333102413476374
17. Crowley, W. R., Rodriguez-Sierra, J. F., & Komisaruk, B. R. (1977). Analgesia induced by vaginal stimulation in rats is apparently independent of a morphine-sensitive process. *Psychopharmacology, 54*(3), 223–225. https://pubmed.ncbi.nlm.nih.gov/413136/
18. American Heart Association. Sexual activity and heart disease (https://www.heart.org/en/health-topics/consumer-healthcare/what-is-cardiovascular-disease/sex-and-heart-disease). Accessed November 9, 2026.
19. Levine, G. N., et al. (2012). Sexual activity and cardiovascular disease: A scientific statement from the American Heart Association. *Circulation, 125*(8), 1058–1072. https://doi.org/10.1161/CIR.0b013e3182447787
20. Valstad, M., Alvares, G. A., Egknud, M., Matziorinis, A. M., Andreassen, O. A., Westlye, L. T., & Quintana, D. S. (2017). The correlation between central and peripheral oxytocin concentrations: A systematic review and meta-analysis. *Neuroscience & Biobehavioral Reviews, 78*, 117–124. https://doi.org/10.1016/j.neubiorev.2017.04.017
21. Feldman, R. (2012). Oxytocin and social affiliation in humans. *Hormones and Behavior, 61*(3), 380–391. https://doi.org/10.1016/j.yhbeh.2012.01.008
22. Bancroft, J., & Janssen, E. (2000). The dual control model of male sexual response: A theoretical approach to centrally mediated erectile dysfunction. *Neuroscience & Biobehavioral Reviews, 24*(5), 571–579. https://doi.org/10.1016/S0149-7634%2800%2900024-5
23. NHS. Antidepressants (https://www.nhs.uk/medicines/antidepressants/). Accessed November 9, 2026.
24. NHS. Low sex drive (https://www.nhs.uk/conditions/loss-of-libido/). Accessed November 9, 2026.
25. NHS. Erection problems (https://www.nhs.uk/conditions/erection-problems-erectile-dysfunction/). Accessed November 9, 2026.
26. Basson, R. (2000). The female sexual response: A different model. *Journal of Sex & Marital Therapy, 26*(1), 51–65. https://doi.org/10.1080/009262300278641
27. Brotto, L. A., & Yule, M. (2017). Asexuality: Sexual orientation, paraphilia, sexual dysfunction, or none of the above? *Archives of Sexual Behavior, 46*(3), 619–627. https://doi.org/10.1007/s10508-016-0802-7
28. Ryan, R. M., & Deci, E. L. (2000). Self-determination theory and the facilitation of intrinsic motivation, social development, and well-being. *American Psychologist, 55*(1), 68–78. https://doi.org/10.1037/0003-066X.55.1.68
29. Maxwell, J. A., Muise, A., MacDonald, G., Day, L. C., Rosen, N. O., & Impett, E. A. (2017). How implicit theories of sexuality shape sexual and relationship well-being. *Journal of Personality and Social Psychology, 112*(2), 238–279. https://doi.org/10.1037/pspi0000078
30. Hazan, C., & Shaver, P. (1987). Romantic love conceptualized as an attachment process. *Journal of Personality and Social Psychology, 52*(3), 511–524. https://doi.org/10.1037/0022-3514.52.3.511

31. Muise, A., & Impett, E. A. (2015). Good, giving, and game: The relationship benefits of communal sexual motivation. *Social Psychological and Personality Science, 6*(2). https://doi.org/10.1177/1948550614553641
32. RAINN. Consent 101 (https://rainn.org/share-the-facts/consent-101-respect-boundaries-and-building-trust/). Accessed November 9, 2026.
33. Dewitte, M., Carvalho, J., Corona, G., Limoncin, E., Pascoal, P., Reisman, Y., & Štulhofer, A. (2020). Sexual desire discrepancy: A position statement of the European Society for Sexual Medicine. *Sexual Medicine, 8*(2), 121–131. https://doi.org/10.1016/j.esxm.2020.02.008
34. Mallory, A. B. (2022). Dimensions of couples' sexual communication, relationship satisfaction, and sexual satisfaction: A meta-analysis. *Journal of Family Psychology, 36*(3), 358–371. https://doi.org/10.1037/fam0000946
35. Muise, A., Impett, E. A., & Desmarais, S. (2013). Getting it on versus getting it over with: Sexual motivation, desire, and satisfaction in intimate bonds. *Personality and Social Psychology Bulletin, 39*(10), 1320–1332. https://doi.org/10.1177/0146167213490963
36. Kim, J. J., Muise, A., Sakaluk, J. K., Rosen, N. O., & Impett, E. A. (2020). When tonight is not the night: Sexual rejection behaviors and satisfaction in romantic relationships. *Personality and Social Psychology Bulletin, 46*(10), 1476–1490. https://doi.org/10.1177/0146167220907469
37. NHS. About vaginal oestrogen (https://www.nhs.uk/medicines/hormone-replacement-therapy-hrt/vaginal-oestrogen/about-vaginal-oestrogen/). Accessed November 9, 2026.
38. Brotto, L. A., et al. (2021). A randomized trial comparing group mindfulness-based cognitive therapy with group supportive sex education and therapy for the treatment of female sexual interest/arousal disorder. *Journal of Consulting and Clinical Psychology, 89*(7), 626–639. https://doi.org/10.1037/ccp0000661
39. National Institute of Mental Health. My mental health: Do I need help? (https://www.nimh.nih.gov/health/publications/my-mental-health-do-i-need-help). Accessed November 9, 2026.
40. Turk, F., & Waller, G. (2020). Is self-compassion relevant to the pathology and treatment of eating and body image concerns? A systematic review and meta-analysis. *Clinical Psychology Review, 79*, 101856. https://doi.org/10.1016/j.cpr.2020.101856
41. Kalmbach, D. A., Arnedt, J. T., Pillai, V., & Ciesla, J. A. (2015). The impact of sleep on female sexual response and behavior: A pilot study. *The Journal of Sexual Medicine, 12*(5), 1221–1232. https://doi.org/10.1111/jsm.12858
42. U.S. Department of Veterans Affairs, National Center for PTSD. PTSD and the family (https://www.ptsd.va.gov/professional/treat/specific/ptsd_family.asp). Accessed November 9, 2026.
43. Huang, S., Li, Z., & Santtila, P. (2024). The effectiveness of online sensate focus exercises in enhancing sexual function and intimacy among Chinese heterosexual couples: A randomized controlled trial. *Journal of Sex & Marital Therapy, 50*(6), 707–724. https://doi.org/10.1080/0092623X.2024.2355229
44. Frühauf, S., Gerger, H., Schmidt, H. M., Munder, T., & Barth, J. (2013). Efficacy of psychological interventions for sexual dysfunction: A systematic review and meta-analysis. *Archives of Sexual Behavior, 42*(6), 915–933. https://doi.org/10.1007/s10508-012-0062-0
45. National Institute of Mental Health. Depression. https://www.nimh.nih.gov/health/publications/depression. Accessed November 9, 2026.

46. National Institute of Mental Health. Generalized anxiety disorder. https://www.nimh.nih.gov/sites/default/files/documents/health/publications/generalized-anxiety-disorder-gad/generalized_anxiety_disorder.pdf. Accessed November 9, 2026.
47. 988 Suicide & Crisis Lifeline. https://988lifeline.org. Accessed November 9, 2026.
48. Centers for Disease Control and Prevention. Guide to taking a sexual history (https://www.cdc.gov/sti/hcp/clinical-guidance/taking-a-sexual-history.html). Accessed November 9, 2026.
49. Centers for Disease Control and Prevention. Getting tested for STIs (https://www.cdc.gov/sti/testing/). Accessed November 9, 2026.
50. Lindau, S. T., Schumm, L. P., Laumann, E. O., Levinson, W., O'Muircheartaigh, C. A., & Waite, L. J. (2007). A study of sexuality and health among older adults in the United States. *New England Journal of Medicine, 357*(8), 762–774. https://doi.org/10.1056/NEJMoa067423
51. Centers for Disease Control and Prevention. Sexual assault and STIs (https://www.cdc.gov/std/treatment-guidelines/sexual-assault-adults.htm) and clinical guidance for HIV post-exposure prophylaxis (https://www.cdc.gov/hivnexus/hcp/pep/). Accessed November 9, 2026.
52. National Domestic Violence Hotline. Should I go to couples therapy with my abusive partner? (https://www.thehotline.org/resources/should-i-go-to-couples-therapy-with-my-abusive-partner/). Accessed November 9, 2026.
53. Meltzer, A. L., Makhanova, A., Hicks, L. L., French, J. E., McNulty, J. K., & Bradbury, T. N. (2017). Quantifying the sexual afterglow: The lingering benefits of sex and their implications for pair-bonded relationships. *Psychological Science, 28*(5), 587–598. https://doi.org/10.1177/0956797617691361
54. U.S. Department of Justice, Office on Violence Against Women. Resources for victims and survivors. https://www.justice.gov/ovw/resources-victims-and-survivors. RAINN. National Sexual Assault Hotline. https://rainn.org/help-and-healing/hotline. Accessed November 9, 2026.